



FULL TITLE OF THE TRIAL

POSNOC - POSitive Sentinel NOde: adjuvant therapy alone versus adjuvant therapy plus Clearance or axillary radiotherapy. A randomised controlled trial of axillary treatment in women with early stage breast cancer who have metastases in one or two sentinel nodes

SHORT TITLE/ACRONYM

POSNOC - A randomised trial of armpit (axilla) treatment for women with early stage breast cancer

Version and Date of Protocol:

Final Version 4.0 27 September 2017

Sponsor: Derby Teaching Hospitals NHS Foundation Trust

Sponsor Protocol Number: RD-5103-001-13

ISRCTN Number: ISRCTN54765244

ClinicalTrials.gov: NCT02401685

Funder	
UK	Health Technology Assessment (HTA), National Institute for Health Research (NIHR)
ANZ	National Health and Medical Research Council (NHMRC)

Trial Website: www.posnoc.co.uk

Protocol authorised by:

Name & Role	Date	Signature
Associate Professor Amit Goyal Chief Investigator	17/01/2018	
Dr Teresa Grieve (Sponsors Representative)	9 th January 2018	

ROLES AND RESPONSIBILITIES

Study personnel

Role	Name & Location	Telephone	E-mail
UK Chief Investigator	Amit Goyal Royal Derby Hospital, Derby, UK	01332785538 07588514469	amit.goyal@nhs.net
ANZ Chief Investigator	Bruce Mann The Victorian Comprehensive Cancer Centre,	+61 (03) 9347-6301	bruce.mann@mh.org.au
Sponsor's Representative	Teresa Grieve Royal Derby Hospital, Derby, UK	01332724710	Teresa.grieve@nhs.net
Statisticians	Alan Montgomery Nottingham CTU	01158231612	Alan.montgomery@nottingham.ac.uk
UK Trial Manager	Shabina Sadiq Nottingham CTU	01158231566	posnoc@nottingham.ac.uk
ANZ Trial Manager	Heath Badger, ANZ Breast Cancer Trials Group	+61 (02) 4925 3022	Posnoc@bctrials.org.au
Radiotherapy Quality Assurance	National Radiotherapy Trials QA (RTTQA) Group	02038262320	posnocqa.enh-tr@nhs.net

Co investigators

Name	Address	Telephone	E-mail
Malcolm W Reed	University of Sussex, Brighton	01273 877575	m.reed@bsms.ac.uk
Lesley Fallowfield	University of Sussex, Brighton	01273873015	l.j.fallowfield@sussex.ac.uk
Valerie Jenkins	University of Sussex, Brighton	01273873016	val@sussex.ac.uk
Laura Ternent	Newcastle University, Newcastle	01912087083	Laura.ternent@newcastle.ac.uk
David Dodwell	St James Hospital Leeds	01132068938	david.dodwell@leedsth.nhs.uk
Robert Coleman	Weston Park Hospital, Sheffield	01142265213	r.e.coleman@sheffield.ac.uk
Ian Ellis	University of Nottingham	01159691169	ian.ellis@nottingham.ac.uk
Lelia Duley	Nottingham CTU	01158231611	lelia.duley@nottingham.ac.uk
Diane Whitham	Nottingham CTU	01158231616	diane.whitham@nottingham.ac.uk
Vaughan Keeley	Royal Derby Hospital, Derby	01332783075	Vaughan.keeley@nhs.net
Patricia Fairbrother	Independent Cancer Patients' Voice		pat@icpv.org.uk

The following groups have endorsed this study:

UK Breast Intergroup

NCRI Breast Clinical Studies Group

Independent Cancer Patients' Voice

Association of Breast Surgery(UK)

NCRI Radiotherapy Clinical Studies Group

Lymphoedema Support Network

UK STUDYCOORDINATING CENTRE

Trial Manager

Nottingham Clinical Trials Unit

Nottingham Health Science Partners

Room 2201, C Floor, South Block
Queens Medical Centre
Nottingham
NG7 2UH

Tel: 01158231566

Fax: 01157484092

Email: posnoc@nottingham.ac.uk

AUSTRALIA AND NEW ZEALAND STUDY COORDINATING CENTRE

Breast Cancer Trials

Level 4

175 Scott St

Newcastle NSW 2300

Australia

Tel: +61 (02) 4925 3022

Fax: +61 (02) 4925 3068

Posnoc@bctrials.org.au

The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.



TABLE OF CONTENTS

ROLES AND RESPONSIBILITIES.....	2
TABLE OF CONTENTS.....	4
SUMMARY.....	6
TRIAL FLOW CHART.....	7
1 BACKGROUND AND RATIONALE.....	8
1.1 Sentinel Node Biopsy.....	8
1.2 Axillary Treatment and Arm Morbidity.....	8
1.3 Adjuvant Therapy.....	8
1.4 Evidence from Systematic Reviews and Randomised Trials.....	9
1.5 Why We Need a Trial Now.....	10
1.6 Axillary ultrasound (AUS) detected nodal disease.....	11
1.7 Extracapsular invasion.....	12
2 OBJECTIVES.....	13
2.1 Primary Objective.....	13
2.2 Secondary Objectives.....	13
3 TRIAL DESIGN.....	14
3.1 Randomisation.....	14
4 PARTICIPANTS.....	15
4.1 Inclusion Criteria.....	15
4.2 Exclusion Criteria.....	15
4.3 Trial Interventions.....	16
5 OUTCOMES.....	17
5.1 Primary Outcome.....	17
5.2 Secondary Outcomes.....	17
5.3 Criteria for Measurement of Study Endpoints.....	17
5.3.1 <i>Description and Definition of Outcomes.....</i>	<i>17</i>
5.3.2 <i>Clinical Care Following Disease Recurrence or Progression.....</i>	<i>19</i>
6 PARTICIPANT IDENTIFICATION.....	20
6.1 Pathway 1: After Primary Breast Surgery.....	21
6.2 Pathway 2: Before Primary Breast Surgery.....	22
6.3 Pathway 3: Post Chemotherapy.....	23
6.4 Pathway 4: Sentinel node biopsy prior to neoadjuvant therapy.....	24
6.5 Registration Procedures.....	25
6.6 Randomisation.....	25
7 TREATMENT PLAN.....	26
7.1 Primary Breast Surgery.....	26
7.1.1 <i>Breast Conserving Surgery.....</i>	<i>26</i>
7.1.2 <i>Mastectomy.....</i>	<i>26</i>
7.2 Sentinel Node Biopsy.....	26
7.3 Pathology.....	26
7.4 Axillary Treatment.....	27
7.4.1 <i>Axillary Node Clearance (ANC) for Standard Care.....</i>	<i>27</i>
7.4.2 <i>Axillary Radiotherapy (ART) for Standard Care.....</i>	<i>27</i>
7.5 Adjuvant Therapy.....	27
7.5.1 <i>Chemotherapy and Endocrine Therapy.....</i>	<i>27</i>
7.5.2 <i>Radiotherapy.....</i>	<i>27</i>
8 STUDY PARAMETERS.....	29
8.1 Study Calendar.....	29
8.2 Study Assessments.....	31
8.2.1 <i>Baseline.....</i>	<i>31</i>
8.2.2 <i>Follow-up Assessments.....</i>	<i>31</i>
9 END OF PARTICIPATION AND TRIAL.....	32
9.1 Participant Retention.....	32
9.3 Trial Discontinuation.....	32
9.4 Definition of End of Study.....	33
10 ADVERSE EVENTS.....	34

10.1	Serious Adverse Events	34
10.2	Safety Reporting.....	35
10.3	Annual Progress Reports	35
11	DATA COLLECTION	36
11.1	Data Forms and Data Entry	36
11.2	Flagging with NHS Digital.....	36
11.3	Source Data	36
11.4	Data Access	37
12	STATISTICS	38
12.1	Sample Size	38
12.1.1	<i>Drop-out Rate</i>	38
12.1.2	<i>Non-inferiority Margin</i>	38
12.2	Statistical Methods	39
12.3	Statistical Analysis.....	39
12.4	Analysis Population	41
12.5	Missing Data	41
13	HEALTH ECONOMIC ANALYSIS	42
14	REGULATORY AND PARTICIPANT INFORMED CONSENT	44
14.1	Research Ethics Approval	44
14.1.1	<i>Protocol Amendments</i>	44
14.1.2	<i>Ethical Considerations</i>	44
14.2	Consent.....	45
14.3	Confidentiality.....	45
14.4	Access to Data	46
15	QUALITY ASSURANCE	47
15.1	Site Start-up and Training.....	47
15.2	Monitoring	47
15.3	Radiotherapy Quality Assurance	48
16	TRIAL MANAGEMENT	50
16.1	Trial Management Group.....	50
16.2	Trial Steering Committee.....	50
16.3	Data Monitoring Committee	50
16.4	Indemnity.....	51
16.5	Public and Patient Involvement	51
16.6	Security and Back-up of Data	51
16.7	Archiving	51
16.8	Dissemination policy.....	52
16.8.1	<i>Dissemination plan</i>	52
16.8.2	<i>Policy for Publication and Authorship</i>	52
APPENDICES		58
1.	GLOSSARY OF ABBREVIATIONS / ACRONYMS.....	58

SUMMARY

POSNOC is a pragmatic, randomised, multicentre, non-inferiority, international (UKANZ) trial.

Aim

For women with early stage breast cancer and one or two nodes with macrometastases, to assess whether adjuvant therapy alone is no worse than adjuvant therapy plus axillary treatment, in terms of axillary recurrence within 5 years.

Patient Population

Women with unifocal or multifocal invasive breast cancer, largest primary lesion ≤ 5 cm, who undergo sentinel node biopsy (SNB) and have 1 or 2 nodes with macrometastases (>2 mm or defined as macrometastasis on molecular assay).

Stratification

- Institution,
- Age (<50 , ≥ 50)
- Breast-conserving surgery (BCS) or mastectomy
- Estrogen receptor (ER) status (positive, negative)
- Number of positive nodes (1, 2)
- Intra-operative sentinel assessment using OSNA (yes, no)

Interventions

The study will compare adjuvant therapy alone with adjuvant therapy plus axillary treatment (axillary node clearance (ANC) or axillary radiotherapy (ART)).

Primary Outcome

Axillary recurrence at 5 years

Secondary Outcomes

- Arm morbidity
- Quality of life
- Anxiety
- Local (breast or chest wall) recurrence
- Regional (nodal) recurrence
- Distant metastasis
- Time to axillary recurrence
- Axillary recurrence free survival
- Disease free survival
- Overall survival
- Contralateral breast cancer
- Non-breast malignancy
- Economic evaluation

Sample Size

1900 participants


Duration of Recruitment and Follow-up

Recruitment will be for 51 months. Participants will be followed up for 5 years.

Adjuvant Therapy

All participants will receive adjuvant systemic therapy (chemotherapy and/or endocrine therapy). All participants may receive breast/chest wall radiotherapy. Axillary and supraclavicular fossa radiotherapy is not allowed when randomised to adjuvant therapy alone.

TRIAL FLOW CHART



1 BACKGROUND AND RATIONALE

1.1 Sentinel Node Biopsy

Each year, more than 48,000 women in the UK are diagnosed with breast cancer and the majority (80%) undergo surgical treatment¹. There is robust evidence that in early breast cancer, sentinel node biopsy (SNB) accurately stages the axilla with low axillary recurrence rates, comparable survival and reduced morbidity when compared with axillary node clearance (ANC)²⁻⁶. SNB has reduced risk of lymphoedema, shoulder discomfort, sensory deficits, and infections than ANC. Quality of life is superior for patients who undergo SNB³. These results have led to widespread adoption of this technique around the world and the Department of Health in the UK supported the NEW START training program which trained UK breast surgeons in this technique⁷.

Currently, it is widely accepted that the patient whose sentinel node (SN) is tumour-free does not require further axillary-specific treatment. A quarter of patients (9,600 patients per year) are found to have SN metastases. These patients return for a second operation, ANC or receive axillary radiotherapy (ART).

1.2 Axillary Treatment and Arm Morbidity

The current UK National Institute of Clinical Excellence (NICE) guidelines⁸ recommend axillary node clearance or axillary radiotherapy for women with early stage breast cancer and one or two sentinel node metastases. This recommendation is based on the assumption that axillary treatment reduces the risk of axillary recurrence, and might improve survival. Axillary node clearance is usually a second operation, but some hospitals use intra-operative sentinel node assessment and so perform axillary node clearance at the same time as breast conserving surgery or mastectomy.

Axillary treatment damages lymphatic drainage from the arm, and women may subsequently develop lymphoedema. As well as the discomfort of arm swelling this causes restricted shoulder movement, pain, numbness and other sensory problems. For example, following axillary treatment 1 in 5 to 1 in 10 people develop lymphoedema, and between 1 in 5 and 1 in 100 have impairment of shoulder function. Sensory changes and pain may occur in up to 1 in 3 patients^{3,9-14}. These adverse effects interfere with daily activities, are distressing, impair quality of life and are costly to the NHS in terms of rehabilitative treatments (such as physiotherapy and lymphoedema clinics), as they are often irreversible and symptom relief is difficult.

1.3 Adjuvant Therapy

Most women with 1 or 2 sentinel node metastases will receive systemic therapy (chemotherapy and/or endocrine therapy). Radiotherapy to the breast is required after breast-conserving surgery and may be given after mastectomy. The aim of axillary treatment (ANC or ART) is to reduce the risk of axillary recurrence. However, systemic adjuvant therapy is now so effective for early breast

cancer that axillary treatment may offer no additional protection against axillary recurrence, and so may be overtreatment. This hypothesis is supported by several small studies¹⁵.

In the past, information from axillary clearance with regard to the number of nodes with cancer was used to guide systemic chemotherapy and hormone therapy. However, decisions about these adjuvant therapies are now more commonly based on biological tumour markers and molecular determinants of prognosis and predictors of treatment benefit. Early data from the EORTC AMAROS trial¹⁶ suggests that once patients are found to have cancer spread to lymph glands by SNB; this information is adequate to guide systemic adjuvant therapy, without the need to remove further nodes. In this study chemotherapy was given to 58% (175 of 300) of patients who had axillary node clearance and 61% (162 of 266) who had axillary radiotherapy; and hormone therapy was given to 78% (235 of 300) who had axillary node clearance and 76% (203 of 266) who had axillary radiotherapy. Patients with cancer spread to four or more nodes are candidates for chest wall and supraclavicular fossa radiotherapy⁸. The proportion of patients having four or more additional positive lymph nodes (besides sentinel node) in the AMAROS trial was low (8%)¹⁷. This figure is estimated to be lower than 5% in the POSNOC trial as pre-operative axillary ultrasound is routinely performed in current practice before sentinel node biopsy and patients with FNA or biopsy proven high volume disease on imaging undergo ANC rather than SNB¹⁸.

1.4 Evidence from Systematic Reviews and Randomised Trials

There is no clear evidence that one form of axillary treatment is better than the other. Axillary radiotherapy has been compared with axillary node clearance in four randomised trials^{11;17;19;20} and several retrospective studies^{21;22}, with no difference in survival. Regional recurrence with both is low at 1% or 2%; however, both have significant morbidity and are costly to the health services^{10;11;13;20-23}. Lymphoedema, sensory changes and pain are more common and more debilitating following axillary node clearance than axillary radiotherapy. Shoulder function impairment is roughly the same after both treatments.

Axillary treatment may now be over treatment for early breast cancer; as diagnosis tends to be earlier so patients present with smaller tumours and a low axillary tumour burden; adjuvant therapy has improved and is better at preventing breast and axillary recurrence²⁴; and sentinel node biopsy has already removed the lymph nodes most likely to have metastasis. For example, in one study less than half of patients with sentinel node metastases had metastasis in the remaining nodes at axillary node clearance⁷. Moreover, if adjuvant therapy includes radiotherapy to the breast or chest wall, the lower axilla may be treated inadvertently as it may be included in the tangential irradiation field, and some lower level axillary nodes may be removed at mastectomy²⁵.

There are three randomised trials assessing axillary treatment. The first¹⁹ was a three arm study that recruited 1079 clinically node-negative women. They were randomised to receive either radical mastectomy (mastectomy with axillary node clearance), or total mastectomy with axillary irradiation, or total mastectomy alone without axillary treatment. Women had larger tumours, higher axillary tumour burden compared with today's patients and they did not routinely receive adjuvant systemic therapy. All three arms had similar 25 year overall survival, suggesting that axillary treatment did not improve survival.

The second study²⁶ randomised 435 clinically node-negative women to breast conservation without axillary treatment or breast conservation plus axillary radiotherapy. Axillary recurrence was low in both groups (no axillary treatment 1.5% vs. 0.5% axillary radiotherapy). Both arms had similar disease free survival.

In the third more recent trial²⁷ patients with tumours less than 5 cm in size, treated by breast conserving surgery and whole breast radiotherapy, with sentinel node metastases, were randomised to axillary node clearance (n=445) or not (n=446). Axillary recurrence was low, and there were no clear differences between the two groups (axillary clearance 0.5% vs. no axillary clearance 0.9%) at 6.3 years. This study has been criticised for not meeting its recruitment goal, approximately 40% of patients had micrometastases, too many patients were lost to follow-up (19.4%), and there was a lack of radiation therapy quality assurance. Missing radiotherapy data for more than two-thirds of all randomised patients, significant number of radiotherapy protocol deviations (18.9% of patients received regional nodal RT using ≥ 3 fields) and the use of high tangent fields²⁸ in half of the patients seriously confound the interpretation of this trial. A recent post Z11 survey shows that most US radiation oncologists treat the undissected axilla in women with micro or macro metastases with axillary radiotherapy²⁹. Therefore, it appears that clinicians in the US are interpreting Z11 in line with AMAROS study that showed that axillary radiotherapy is a less morbid alternative to axillary lymph node dissection (ALND) in women with low volume nodal disease³⁰. This is not surprising given that Z11 lacked radiotherapy quality assurance. Furthermore, the generalisability of the results is limited mastectomy patients were excluded and pre-operative axillary ultrasound was not performed in contrast to standard NHS practice.

A recent meta-analysis¹⁵ of randomised trials and observational studies which included patients who had sentinel node biopsy concluded that more evidence is needed to guide management of the axilla in patients with early breast cancer and sentinel nodes metastasis.

1.5 Why We Need a Trial Now

Axillary management consensus meeting (Association of Breast Surgery Conference, Bournemouth, 21 May, 2012) vote of 226 UK consultant breast surgeons from 155 hospitals, showed that clinicians

are currently at equipoise in terms of effective axillary treatment. Voting results showed that 68.2% would randomise patients in the POSNOC trial, 11.7% were unsure, while 20.1% would not. NICE guidelines stress the need for research to clarify optimum treatment of the axilla in patients with sentinel node metastases⁸. The POSNOC trial will provide evidence relevant to patients and to the NHS. The protocol has been designed to integrate into current NHS practice.

Biological factors may be more important for recurrence than surgical removal or radiation eradication of axillary nodes. If axillary surgery is merely a staging or diagnostic procedure, then adverse effects are likely to be minimal if it is omitted and sentinel node biopsy alone is used to guide subsequent treatment in women with early stage breast cancer who have 1 or 2 nodes with metastases. Also, axillary treatment (axillary node clearance or axillary radiotherapy) was introduced several decades ago without formal evaluation and is associated with significant short- and long-term morbidity. Since axillary treatment was introduced, chemotherapy and hormone therapy have dramatically improved outcome. Therefore, it is timely to assess whether adjuvant therapy alone is an acceptable alternative to adjuvant therapy plus axillary treatment.

The hypothesis of the POSNOC trial is that low axillary tumour burden patients with macrometastases in 1 or 2 SNs, receiving systemic therapy, would have non-inferior outcomes whether they are randomised to adjuvant therapy alone or adjuvant therapy plus axillary treatment (ANC or ART).

1.6 Axillary ultrasound (AUS) detected nodal disease

AUS has been embedded into routine breast practice in the UK. The sensitivity of axillary ultrasonography (AUS) has increased in recent years, enabling detection of even low-volume axillary nodal metastases. Women with ultrasound-detected disease undergo ANC, as they are believed to have a higher axillary tumour burden. Studies show that overall, patients with AUS-positive disease have a larger number of involved nodes and are more likely to have tumours with extracapsular invasion at ANC³¹⁻³⁵. Proceeding to ANC in patients with axillary metastasis detected by AUS therefore appears to be the logical step in management. However, ANC cannot be expected to benefit around 40 percent of patients in the AUS-positive group with only one or two nodes with macrometastases at ANC³³⁻³⁵. Our data shows that tumour size and number of abnormal nodes seen on ultrasound scan could potentially be used to select patients who could proceed to SNB instead of ANC. As an example, a woman with tumour size of 20 mm or less and one abnormal node on ultrasound scan would have a 78 per cent chance of having two or fewer metastases at ANC³⁶. Therefore, women with ultrasound detected disease who proceed to sentinel node biopsy and are found to have 1 or 2 nodes with macrometastases, will be eligible for POSNOC.

1.7 Extracapsular invasion

Most of the published literature relating to extracapsular invasion (ECI) in nodes removed during sentinel node biopsy is from pre-axillary ultrasound era. AUS is able to detect most women with macroscopic ECI and these women undergo ANC rather than SNB. Women found to have ECI during SNB are more likely to have microscopic disease and role of ANC in these women is unclear. Some centres in the UK use intraoperative molecular tests (e.g. OSNA) to analyse the SN. During this test, the whole lymph node is mashed by the technician and therefore the presence or absence of ECI is not known. To ensure consistency between OSNA and non-OSNA centres, and to clarify the role of axillary treatment in women with microscopic ECI, these women will be eligible for POSNOC.

2 OBJECTIVES

2.1 Primary Objective

For women with early stage breast cancer and one or two nodes with macrometastases, to assess whether adjuvant therapy alone is no worse than adjuvant therapy plus axillary treatment (i.e. less than 2% absolute risk difference), in terms of axillary recurrence within 5 years.

2.2 Secondary Objectives

(i) To assess lymphoedema and other arm morbidity, quality of life, and anxiety at three years for women in the two allocated groups.

(ii) To assess the following outcomes at five years in the two allocated groups:

- local (breast or chest wall) recurrence,
- regional (nodal) recurrence,
- distant metastasis,
- time to axillary recurrence,
- survival (axillary recurrence free, disease free and overall),
- contralateral breast cancer
- non-breast malignancy

(iii) Economic Evaluation

- a. To compare quality-adjusted life years (QALYs) using the area under the curve method based upon responses to the EQ-5D-5L measured at baseline, 3, 6, 12, 24, and 36 months post randomisation.
- b. To compare the cost-effectiveness measured in terms of the incremental cost per reduction in axillary recurrence from the perspective of the NHS over 36 months
- c. To compare the cost-utility based on incremental cost per QALY gained from the perspective of the NHS over 36 months

3 TRIAL DESIGN

POSNOC is a pragmatic randomised, multi-centre, non-inferiority, international (UK-ANZ) trial. Estimated sample size is 1900 participants, with follow up for five years.

3.1 Randomisation

Participants will be allocated to one of the two intervention groups, via a remote, secure internet-based randomisation system developed and maintained by the NCTU.

A nominated member from the local recruiting team will login using their unique username and a password. Once logged in they will enter required participant details and the system will allocate the participant to a treatment arm (in real-time).

Allocations will be assigned using the Simon and Pocock minimisation method (with an 80% chance of attempting to balance), in order achieve a 1:1 ratio. The minimisation will be stratified by the recruitment site and minimised across the following variables:

- Age (<50, ≥50)
- Breast-conserving surgery (BCS) or mastectomy
- Estrogen receptor status (ER) (positive, negative)
- Number of positive nodes (1, 2)
- Intra-operative sentinel assessment using OSNA (yes, no)

The sequence of treatment allocations will be kept concealed until all interventions have been assigned, recruitment, data collection, and all other trial-related assessments are complete.

It will not be possible to blind clinicians or participants to the treatment allocation. To minimise the potential for bias, we will train and monitor to ensure any breast or chest wall radiotherapy is not influenced by knowledge of whether the woman had axillary treatment, or not. For women randomised intra-operatively we will also monitor to ensure that women allocated no axillary treatment do not have any lymph nodes removed after the sentinel node biopsy. To reduce the potential for participants' expectations to influence their responses to the questionnaires, the participant information leaflet and DVD will emphasise that we do not know whether axillary treatment is worthwhile, and that is why we are doing the study.

A procedure for unblinding is not necessary, as the clinicians and participants will know the study allocation.

4 PARTICIPANTS

Women with unifocal or multifocal invasive breast cancer with the largest lesion ≤ 5 cm, who have undergone sentinel node biopsy, and have 1 or 2 nodes with macrometastases (>2 mm) The criteria and procedure for sampling lymph nodes showing abnormal morphology on ultrasound should be agreed locally. Women will be enrolled from 50 or more academic and non-academic hospitals with an established and functional Breast MDT (multidisciplinary team).

Women will usually be screened for eligibility after their initial breast surgery and sentinel node biopsy. In hospitals where sentinel node histology is checked intra-operatively, allowing axillary treatment if necessary, to be performed at the same time as the breast surgery, women will consent pre-operatively and then randomised intra-operatively if their sentinel nodes are found to be positive. Alternatively these hospitals may follow the non-intraoperative pathway, however women need to be consented at the appropriate time points.

4.1 Inclusion Criteria

Women will be eligible for inclusion only if **ALL** of the following criteria apply:

- 18 years or older
- Unifocal or multi-focal invasive tumour with lesion ≤ 5 cm in its largest dimension, measured pathologically or largest invasive tumour diameter on radiology should be used for women who are randomised intra-operatively or undergo sentinel node biopsy before neoadjuvant therapy (tumour size should be based only on the single largest tumour; do not add the sizes together from the multiple foci)
- At sentinel node biopsy have 1 or 2 nodes with macrometastases (tumour deposit >2.0 mm in largest dimension or defined as macrometastasis on molecular assay)
- Fit for axillary treatment and adjuvant therapy
- Have given written informed consent

4.2 Exclusion Criteria

Women will be excluded if they have:

- Bilateral invasive breast cancer
- more than 2 nodes with macrometastases
- neoadjuvant therapy for breast cancer except
 - if sentinel node biopsy performed prior to neoadjuvant therapy in women with early breast cancer
 - short duration of neoadjuvant endocrine therapy is acceptable (up to 3 months)
- previous axillary surgery on the same body side as the scheduled sentinel node biopsy
- not receiving adjuvant systemic therapy
- previous cancer less than 5 years previously or concomitant malignancy except
 - basal or squamous cell carcinoma of the skin
 - *in situ* carcinoma of the cervix
 - *in situ* melanoma
 - contra- or ipsilateral *in situ* breast cancer

4.3 Trial Interventions

All participants will receive adjuvant therapy according to local guidelines. Adjuvant therapy will include chemotherapy and/or endocrine therapy for all women, and radiotherapy to breast or chest wall if indicated. Human epidermal growth factor receptor 2 (HER2) targeted treatment may also be administered when indicated.

The trial interventions are either:

(i) Adjuvant therapy alone (intervention)

See above for adjuvant therapy. Axillary and supraclavicular fossa radiotherapy is not allowed when randomised to this group.

(ii) Adjuvant therapy *plus* axillary treatment (standard care)

See above for adjuvant therapy. Axillary treatment can be axillary node clearance or axillary radiotherapy as per local guidelines.

5 OUTCOMES

5.1 Primary Outcome

The primary outcome is axillary recurrence at 5 years.

5.2 Secondary Outcomes

Secondary outcomes assessed at 3, 6, 12, 24 and 36 months are:

- Arm morbidity
- Quality of life
- Anxiety
- Economic Evaluation
 - To compare the total average costs of the intervention and standard care
 - To compare QALYs using the area under the curve method based upon responses to the EQ-5D-5L measured at baseline, 3, 6, 12, 24, and 36 months post randomisation.
 - To compare the cost-effectiveness measured in terms of the incremental cost per reduction in axillary recurrence from the perspective of the NHS over 36 months
 - To compare the cost-utility based on incremental cost per QALY gained from the perspective of the NHS over 36 months

Secondary outcomes assessed at 6, 12, 24, 36, 48 and 60 months are:

- Local (breast or chest wall) recurrence
- Regional (nodal) recurrence
- Distant metastasis
- Time to axillary recurrence
- Axillary recurrence free survival
- Disease free survival
- Overall survival
- Contralateral breast cancer
- Non-breast malignancy

5.3 Criteria for Measurement of Study Endpoints

5.3.1 Description and Definition of Outcomes

- **Axillary recurrence** is defined as pathologically (cytology or biopsy) confirmed recurrence in lymph nodes draining the primary tumour site, i.e. nodes in the ipsilateral axilla, infraclavicular fossa, supraclavicular fossa and interpectoral area.
- **Arm morbidity** will be assessed with questionnaires completed by participants during clinical or telephone follow-up - a) the definition of lymphoedema is 'yes' to the two questions

participants will be asked-'heaviness during the past year' and 'swelling now'. These two questions come from the Lymphoedema and Breast Cancer Questionnaire (LBCQ)³⁷; b) Shoulder, arm and hand disability is defined as a change from baseline in the QuickDASH³⁸ score of at least 14 points.

Quality of life will be assessed with the Functional Assessment of Cancer Therapy-Breast+4 (FACT B+4) questionnaire³⁹ which asks about physical, social, emotional, functional well-being as well as breast cancer concerns and arm morbidity. Participants indicate, using a five-point scale ranging from 0 (not at all), 1 (a little bit), 2 (somewhat), 3 (quite a bit), to 4 (very much), to what degree each item has applied over the past 7 days. The scores of negatively framed statements are reversed for analysis. High FACT scores equate with a good quality of life and lower scores with a poorer one.

The primary Quality of life endpoint is the FACT-B+4 trial outcome index (TOI) score (The TOI score is a sum of the scores of the 28 items included in the physical well-being (7 items), functional well-being (7 items) and breast cancer subscales(14 items) of the FACT-B+4 (range from 0-112)). A change of at least 5 points from baseline in the TOI, is considered to be clinically relevant minimally important difference⁴⁰.

The total FACT B+4 score reflects global quality of life and comprises the physical (7 items), functional (7 items), social (7 items) and emotional well-being (6 items) plus the breast cancer concerns and arm morbidity scale (14 items) (total of 41 items; total score ranges from 0-164). The 5 item arm morbidity subscale score comprises the sum of the scores from items B3, B10, B11, B12, B13 (range 0-20) and will be analysed separately. Analyses will include the proportion of participants in the two allocated treatment groups reporting 'somewhat', 'quite a bit' and 'very much' for each of the 5 items.

- **Anxiety** will be assessed using the Spielberger State/Trait Anxiety Inventory⁴¹. The STAI consists of 2 questionnaires with 20 items. It assesses anxiety proneness (Trait) and the current state of anxiety or anxiety change (State). Each item is rated on a four point Likert scale. High STAI scores signify greater anxiety. The Trait anxiety is measured only once and the State at each time point.
- **Economic outcomes** will use the EQ-5D-5L utility scores as a measure of health outcome. For more details, see section 13.
- **Local (breast or chest wall) recurrence** is defined as pathologically (cytology or biopsy) confirmed recurrence after mastectomy in the skin or soft tissue of the chest wall within the anatomical area bounded by the mid-sternal line, the clavicle, the posterior axillary line and the costal margin or any type of breast carcinoma in the breast after conservation therapy. The date of local recurrence is the date on which the pathology report confirms recurrence.
- **Regional (nodal) recurrence** is defined as pathologically (cytology or biopsy) confirmed recurrent tumour in the lymph nodes in the ipsilateral axilla, infraclavicular, supraclavicular fossa,

interpectoral area or ipsilateral internal mammary chain. The date of regional recurrence is the date on which the pathology report confirms recurrence.

- **Distant metastasis** is defined as confirmed metastasis (positive pathology and/or definitive evidence on imaging) in all other sites of recurrence and may include those classified as: soft-tissue category, visceral category, central nervous system and skeletal spread. The date of distant metastasis is the date on which imaging or pathology report (whichever comes first) confirms metastasis.
- **Time to axillary recurrence** is the time between the date of randomisation and the date of axillary recurrence, measured in days. The date of axillary recurrence is the date on which pathology report confirms recurrence.
- **Axillary recurrence free survival** is the time between the date of randomisation and date of confirmed axillary recurrence or date of death, whichever comes first, measured in days. Participants who did not experience axillary recurrence and are still alive will be censored at the date of last follow up.
- **Disease free survival** is defined as the time between the date of randomisation and the date of disease progression (i.e. local or regional recurrence or distant metastasis) or death, whichever comes first, measured in days. Participants who do not have progression and are still alive will be censored at the date of last follow up.
- **Overall survival** is the time between the date of randomisation and the date of death from any cause. Participants who are still alive will be censored at the date of last follow up.
- **Contralateral breast cancer** is new primary malignancy in the opposite breast unless obviously contiguous with recurrent chest wall disease or proven on cytology/biopsy to be of metastatic origin.
- **Non-breast cancer** is any new non-breast primary malignancy, except for adequately treated, superficial squamous or basal cell carcinoma of the skin, or carcinoma in situ of the cervix.

5.3.2 Clinical Care Following Disease Recurrence or Progression

If there is disease recurrence and progression then participants should be treated according to local guidelines.

6 PARTICIPANT IDENTIFICATION

Potential trial participants will be identified at the routine multi-disciplinary meetings. The trial will be introduced and discussed with the patient by the treating clinician during the results clinic appointment (depending on how the patient reacts to the results) followed by a further discussion with the research nurse. Patient information leaflet (PIL) will be given to the patient. Patient information DVD will be used as an adjunct to PIL. The DVD will contain trial explanation together with frequently asked questions relating to randomisation and study visits. If time allows, the DVD will be viewed by the woman in the presence of the research nurse. Women will be encouraged to take the information home and discuss the trial with their family ahead of making an informed decision. Patient will be given at least one day to consider the information and reach a decision, this may include coming for another clinic visit.

Based on local hospital practices, there are two participant pathways for recruitment:

(i) *After primary breast surgery*: In most hospitals results of the sentinel node biopsy are not available immediately. Women who have a positive sentinel node will be approached with regard to participation in the trial at the first post-operative clinic visit (See section 6.1).


(ii) *Before primary breast surgery*: In a few hospitals (less than 10%) sentinel node assessment is performed in the operating room during primary breast surgery. In these hospitals, women will be approached with regard to the study before surgery. If they are willing to participate consent will be obtained prior to surgery, but they will be randomised intra-operatively only if the sentinel node/s are confirmed as positive (See section 6.2).

(iii) *Post chemotherapy*: Some patients may be psychologically overwhelmed by their post-surgery results and therefore timing to discuss the study needs to be tailored to a patient's condition. Women may be approached to discuss the trial during and after chemotherapy but before starting axillary treatment (ANC or ART) (See section 6.3).

(iv) *Sentinel node biopsy prior to neoadjuvant therapy*: In hospitals where the sentinel node biopsy is performed prior to neoadjuvant therapy patients can be approached during or after their neoadjuvant therapy but before starting axillary treatment (ANC or ART) (See section 6.4).


6.1 Pathway 1: After Primary Breast Surgery

After primary breast surgery and sentinel node biopsy, the clinical team will decide whether the participant would be appropriate for axillary treatment. Eligible women will follow the pathway below:




6.2 Pathway 2: Before Primary Breast Surgery

Eligible women will be consented and registered before primary breast surgery. Women will be randomised intra-operatively if macrometastases are found in 1 or 2 sentinel nodes as shown below:




6.3 Pathway 3: Post Chemotherapy

Eligible women will be consented and registered during or after adjuvant chemotherapy but before starting axillary treatment (ANC or ART). Eligible women will follow the pathway below:



6.4 Pathway 4: Sentinel node biopsy prior to neoadjuvant therapy

Eligible women will be consented and registered before, during or after neoadjuvant therapy but before starting axillary treatment (ANC or ART). Eligible women will follow the pathway below:



6.5 Registration Procedures

1. Confirm eligibility
2. Obtain written informed consent
3. Trial registration
4. Obtain baseline data for demographics, tumour and nodal characteristics, receptor status, arm morbidity, quality of life and anxiety via questionnaires

6.6 Randomisation

The actual allocation will be disclosed on the computer screen and a confirmation email will follow to the person recruiting the participant. For more details, see section 3.1.

7 TREATMENT PLAN

7.1 Primary Breast Surgery

Primary breast surgery can consist of either breast conserving surgery or mastectomy.

7.1.1 Breast Conserving Surgery

Participants will undergo resection of the breast tissue, which contains the primary breast tumour with a clear margin of normal tissue around the periphery of the tumour. Circumferential margins must be assessed and deemed tumour free (as per local protocol) by the institutional pathologist. Participants with involved margins should undergo re-excision or mastectomy as per local protocol. This can be performed before or after randomisation.

7.1.2 Mastectomy

Participants may undergo simple, skin sparing or nipple sparing mastectomy with or without immediate breast reconstruction.

7.2 Sentinel Node Biopsy

Sentinel node biopsy should be performed as per local protocol.. This protocol does not dictate the injection technique or tracer/s to be used, but it is recommended that no more than four gross nodes should be removed. This is not an exclusion criteria. At times, the pathologist may identify more lymph nodes than the surgeon and this is acceptable. For hospitals using intra-operative sentinel node assessment, further sampling after randomisation is not allowed. A blue node is defined as any node which is visibly blue or contiguous with a blue lymphatic vessel. A hot node is defined as any node with an isotope count at least 10% of the hottest node or a node with counts more than 10 times the background count. Any clinically suspicious nodes should also be removed. All nodes should be labelled and sent to the laboratory separately. Participants with failed localisation, undergoing axillary node sampling are eligible for the trial.

The number of nodes removed at sentinel node biopsy will be monitored by reviewing the pathology report. Prompt action will be taken if there is any indication that the protocol is not being adhered to. It is expected that in most cases, the trial management group (TMG) will work with the investigator to improve performance. However, the TMG shall suspend site participation, if necessary.

7.3 Pathology

All lymph nodes should be examined according to the predefined local practice: For pathological examination, each sentinel node is processed separately. Immunohistochemical staining may be used selectively to characterize suspicious or micrometastatic disease. The pathology report should mention the results of H&E/IHC staining for each sentinel node and non-sentinel nodes. In case of additional axillary lymphadenectomy the conclusion should consist of the total number of lymph nodes examined and the number with metastases. Whether the lymph node metastasis is

considered to be a macrometastasis (> 2 mm) or not, and extranodal invasion should be noted in the pathological report. The extent of extranodal invasion (radial extension) should be noted (≤ 2 mm, >2mm).

Intra-operative molecular analyses can be used to analyse either the whole sentinel node or half of the sentinel node with follow-up histopathology on the remaining half to confirm the results according to local guidelines.

7.4 Axillary Treatment

Participants randomised to standard care will undergo axillary node clearance or axillary radiotherapy as per local guidelines.

7.4.1 Axillary Node Clearance (ANC) for Standard Care

Participants will undergo removal of at least level I and II axillary lymph nodes.

7.4.2 Axillary Radiotherapy (ART) for Standard Care

Participants will undergo axillary irradiation therapy as per local radiotherapy guidelines. Radiotherapy will be monitored as described in the accompanying RT planning and delivery guidelines.

7.5 Adjuvant Therapy

7.5.1 Chemotherapy and Endocrine Therapy

All participants will receive currently accepted adjuvant chemotherapy and HER2 targeted treatment according to pre-defined local guidelines. They may receive currently accepted endocrine therapy in addition to chemotherapy or as a single agent endocrine adjuvant treatment.

7.5.2 Radiotherapy

All participants receiving radiotherapy should be treated according to local guidelines and monitored for compliance according to the POSNOC radiotherapy planning and delivery guidelines (see accompanying radiotherapy manual). Accepted fractionations include 40Gy in 15, 50Gy in 25 or 45Gy in 20 daily fractions. All participants should be CT scanned and 3-D planned in order to optimise target coverage and reduce dose to organs at risk. It is not compulsory to outline any target volumes or organs at risk, however if a centre wishes to do so this is encouraged.

Participants in both groups may receive adjuvant breast or chest wall irradiation therapy as per pre-defined local guidelines. The irradiation therapy for this protocol specifically excludes axillary and supraclavicular fossa irradiation when randomized to adjuvant therapy alone as this would confound the issue being addressed by this study. It is recognised that a variable amount of the axilla may be

irradiated unintentionally by standard breast/chest wall tangential fields. However, unless definitive RT to the axilla is selected in preference to axillary node clearance surgery, no attempt should be made to irradiate the axilla by adjustment of the superior/posterior tangential field margins.

8 STUDY PARAMETERS

8.1 Study Calendar

Time point	Intervention/Procedure						
	Pathway 1 After Primary Breast Surgery		Pathway 2 Before Primary Breast Surgery		Pathway 3 Post Chemotherapy		Pathway 4 Post Neoadjuvant therapy
Baseline	<ul style="list-style-type: none"> Verify eligibility Obtain informed consent Trial Registration 		<ul style="list-style-type: none"> Verify eligibility except for sentinel node status, this to be obtained during surgery Obtain informed consent Trial Registration 		<ul style="list-style-type: none"> Verify eligibility Obtain informed consent Trial Registration 		<ul style="list-style-type: none"> Verify eligibility Obtain informed consent Trial Registration
Baseline recordings	Record demographics						
	Record tumour and nodal characteristics Record receptor status				Record tumour and nodal characteristics Record receptor status		Record tumour and nodal characteristics Record receptor status
Baseline questionnaires	<ol style="list-style-type: none"> Lymphoedema and Breast Cancer Questionnaire and QuickDASH Functional Assessment of Cancer Therapy –Breast +4 EQ-5D-5L Spielberger State/Trait Anxiety Inventory 						
Randomisation	Intervention – adjuvant therapy alone Standard Care – adjuvant therapy plus axillary treatment		Intra-operative randomisation following confirmation of eligibility* Intervention – adjuvant therapy alone Standard Care – adjuvant therapy plus axillary treatment		Intervention – adjuvant therapy alone Standard Care – adjuvant therapy plus axillary treatment		Intervention – adjuvant therapy alone Standard Care – adjuvant therapy plus axillary treatment
Baseline recordings			Record tumour and nodal characteristics Record receptor status				
Systemic therapy	Chemotherapy and/or endocrine therapy				Endocrine therapy		
Radiotherapy	Breast or chest wall radiotherapy, if indicated. <i>NB: Axillary and Supraclavicular fossa (SCF) radiotherapy is not allowed in the Intervention group.</i>						
Follow-up (months)	3	6*	12*	24*	36**	48*	60**
Lymphoedema and Breast Cancer Questionnaire and QuickDASH (arm morbidity)		X ^{\$\$}	X ^{\$\$}	X ^{\$\$}	X ^{\$\$}		
Functional Assessment of Cancer Therapy –Breast +4 and	X ^{\$}	X ^{\$}	X ^{\$}	X ^{\$}	X ^{\$}		

EQ-5D-5L (Quality of life)							
Spielberger State/Trait Anxiety Inventory (anxiety)	X\$	X\$	X\$	X\$	X\$		
Assessment of disease recurrence or progression and non-hospital resource use		X	X	X	X	X	X
¥ For ineligible participants, no further data will be collected * Follow up will be clinic visits or telephone consultation (doctor or nurse led) ** Follow up will be clinic visits (doctor or nurse led). \$ Questionnaires are posted by SHORE-C with up to two reminders (by post or telephone). \$\$ Questionnaires are completed in clinic or over telephone by site staff. Shaded areas indicate that those recordings are not performed at that time points in that particular pathway							

NB: the date of the follow-up appointments are calculated from the date of randomisation.

8.2 Study Assessments

8.2.1 Baseline

Baseline data will be collected from the participant's medical notes and questionnaires.

8.2.2 Follow-up Assessments

Follow-up will be at 6, 12, 24, 36, 48 and 60 months post randomisation. The 36 and 60 months follow-up will be a clinic visit (doctor or nurse led), however follow-up at other time points may be conducted by a research nurse over the telephone, according to local practice.

Assessments:

1. Follow-up eCRFs will be completed in the clinic or over the telephone to record information on primary and secondary outcomes at 6, 12, 24, 36, 48 and 60 months,
2. Lymphoedema and Breast Cancer Questionnaire (two questions) and QuickDASH questionnaire will be completed in the clinic or over the telephone at 6, 12, 24 and 36 months,
3. The number of breast cancer related GP visits, physiotherapy visits, lymphoedema visits, pain team visits and hospital admissions will be collected in the clinic or over the telephone at 6,12, 24, 36 and 60 months,
4. Postal questionnaires with up to two reminders, FACT B+4, Spielberger State/Trait Anxiety Inventory and EQ-5D-5L at 3, 6, 12, 24 and 36 months.

9 End of Participation and Trial

9.1 Participant Retention

Once a woman has been randomised, the study site will make every reasonable effort to follow the participant for the entire study duration. We anticipate a maximum of 5% loss to follow up over 5 years.

All reasonable attempts will be made to contact any participant lost to follow-up during the course of the study in order to complete assessments.

9.2 Participant Withdrawal

Women are free to stop study participation at any time without giving a reason. Participants may stop taking part in the Quality of Life study but may continue with the treatment follow-up phase. Stopping participation in the Quality of Life (QoL) study will be recorded in the QoL specific database.

Ceasing participation in the trial follow-up will be recorded in the eCRF. Data already collected will be included in the analysis, unless participants specifically request for their data not to be used.

Women who consent to take part but cease participation prior to randomisation will receive routine NHS care. Further research data will not be collected for these women and any data already collected will be securely kept at site until the end of the study and then destroyed in a confidential manner.

Participants who withdraw will not be replaced.

9.3 Trial Discontinuation

The sponsor and Funder reserve the right to discontinue this study at any time for failure to meet expected recruitment goals, for safety or any other administrative reason. The Sponsor and Funder shall take advice from the Trial Steering Committee as appropriate in making this decision. An independent Data Monitoring Committee shall monitor accumulating data and oversee safety issues. The DMC will advise the TSC if, in its view, there are any ethical or safety issues that may necessitate closure of the trial. For more details about the role of DMC, see section 16.3.

All participation may be stopped if the study sponsor or REC terminate the study prior to the planned end date.

9.4 Definition of End of Study

The end of the study is defined as the “last data collection for the last participant”.

10 Adverse Events

In this study the participants are women with early stage breast cancer, who are at risk of adverse outcome due to their condition. The study is comparing two different policies for treatment of early stage breast cancer, to assess whether their effects are comparable. Adverse events (AE) that could be influenced by the trial interventions are therefore outcomes for the study. Data on these events will be recorded on the case report forms.

10.1 Serious Adverse Events

In accordance with the ICH E2A guideline, a serious adverse event or reaction is any untoward medical occurrence that:

- results in death,
- is life-threatening (i.e. the patient was at risk of death at the time of the event/reaction),
- requires inpatient hospitalisation or results in prolongation of existing hospitalisation,
- results in persistent or significant disability/incapacity,
- is a congenital anomaly/birth defect,
- is a medically important event or reaction (i.e. important medical event/reaction that might not be immediately life-threatening or result in death or hospitalisation but might jeopardise the patient or might require intervention to prevent one of the other outcomes listed in the definition above).

For the POSNOC trial, study treatment consists of standard adjuvant therapy with or without standard axillary treatment. This protocol does not contain investigational agent(s).

For this trial events related to the natural course of the disease are not required to be reported as these are outcomes and will be captured in the CRF.

A SAE will be unexpected if it is not described in this study protocol or recorded as an outcome. All unexpected SAEs will be reported. The PI shall assess expectedness, seriousness and causality. All SAEs will be followed up until the event is considered resolved, or resolved with sequelae.

The assessment of causality will be made by the PI using the following definitions:

- *Definitely Related*: The adverse event is clearly related to the investigational agent/procedure - i.e. an event that follows a reasonable temporal sequence from administration of the study intervention, follows a known or expected response pattern to the suspected intervention, that is confirmed by improvement on stopping and reappearance of the event on repeated exposure and that could not be reasonably explained by the known characteristics of the subject's clinical state.

- *Possibly Related:* An adverse event that follows a reasonable temporal sequence from administration of the study intervention follows a known or expected response pattern to the suspected intervention, but that could readily have been produced by a number of other factors.
- *Not Related:* The adverse event is clearly not related to the investigational agent/procedure- i.e. another cause of the event is most plausible; and/or a clinically plausible temporal sequence is inconsistent with the onset of the event and the study intervention and/or a causal relationship is considered biologically implausible.

10.2 Safety Reporting

Whilst it is not anticipated there will be any serious adverse events directly related to the study, it is important that this protocol includes a process for dealing with any unexpected serious adverse events in the unlikely event they occur.

All unexpected Serious Adverse Events (SAEs) that occur in participants recruited to the POSNOC trial should be reported via the trial SAE form and submitted to Nottingham CTU on fax number 0115 7484091 or nctu-sae@nottingham.ac.uk

PI's have the responsibility of safety reporting for the participants recruited to POSNOC at their sites. It is the responsibility of the CI to review any SAEs submitted by investigator sites.

All unexpected SAEs will be reported to the NCTU within one working day. The Chief Investigator will submit, regularly for the duration of recruitment to the trial or upon request, a safety report to the Data Monitoring Committee which will include all reported unexpected SAEs.

10.3 Annual Progress Reports

The CI will send the annual progress report to the main REC (the anniversary date is the date of the REC 'favourable opinion' letter) and to the sponsor.

11 DATA COLLECTION

The trial data will be collected via the study electronic case report form (eCRF), worksheets and questionnaires. The agreed data collection time points are specified in the Study Calendar in section 8.1.

11.1 Data Forms and Data Entry

Trial data will be recorded prior to discharge from hospital and at each follow up visit. The data will be entered at site into electronic case report forms (eCRF) directly or onto worksheets by study personnel. The worksheet data must be transferred into the eCRF within 5 days of data collection. Participants will be identified only by their unique study number.

The trial database will be developed and maintained by the trial coordinating centre at the Nottingham Clinical Trials Unit (NCTU). Access to the database will be restricted and secure. Data quality and compliance with the protocol will be assessed throughout the trial by central statistical monitoring and site visits.

Processing of trial data and monitoring for consistency, validity and quality will be undertaken by NCTU according to their agreed monitoring plan. Data checks will include out-of-range data, cross-checks for conflicting data, missing data and data queries.

The completed Quality of Life (QoL) questionnaires (FACT B+4, and Spielberger State/Trait Anxiety Inventory) and the EQ-5D-5L will be sent to SHORE-C (Sussex University) for entry into their trial specific questionnaire database. SHORE-C will process and monitor follow-up questionnaire data as the questionnaires are received. The SHORE-C database will be transferred to the NCTU for reconciliation with the NCTU trial database for the purpose of statistical analysis. Participants who do not return their questionnaires will be telephoned after two weeks by SHORE-C researchers. Missing data and data queries will be addressed by SHORE-C directly to the participant by sending a data query by post or telephone.

11.2 Flagging with NHS Digital

All women recruited to the study will be 'flagged' after discharge through the Data Linkage and Extract Service at NHS Digital. Information from NHS Digital may be used to help contact participants, check their health status. This will avoid causing unnecessary distress to the family, as it will avert inappropriate contact if the woman has died following discharge from hospital. It will reduce losses to follow up and enable us to report extended 10 year follow-up data of the study.

11.3 Source Data

Source documents shall be filed at the local site and may include but are not limited to consent forms, current medical records, worksheets and participant questionnaires. QoL and EQ-5D-5L

questionnaires will be held at SHORE-C. For some data the eCRF's may be the primary source document data.

11.4 Data Access

Access to the trial datasets will be limited to the central co-ordinating site (NCTU) and to the trial statisticians. The datasets will be provided to the sponsor at the end of the trial for archiving purposes.

12 STATISTICS

The trial statisticians will draft the Statistical Analysis Plan (SAP), which will be reviewed by the Trial Management Group (TMG), the Trial Steering Committee (TSC), and the Data Monitoring Committee (DMC). The finalised SAP will be approved and signed by the CI, trial statistician and chair of the TSC.

12.1 Sample Size

For women who have adjuvant therapy plus axillary treatment, 2% have axillary recurrence at 5 years⁴².

With an assumed axillary recurrence rate in the adjuvant therapy alone group also of 2%, absolute non-inferiority margin of 2%, 80% power and 2.5% one-sided alpha, a total of 1700 women are required for analysis in order to show that the axillary recurrence rate in the adjuvant therapy alone group at 5 years is not more than 4% using the Miettinen-Nurminen score interval⁴³. Allowing for 10% non-compliance with treatment allocation and non-collection of primary outcome data, that target sample size to be randomised is 1900.

12.1.1 Drop-out Rate

Within the NHS, failure to attend for breast cancer follow up clinic visits is unusual. For this study, we anticipate a maximum of 5% loss to follow up over 5 years. This is realistic, for example in one recent UK breast cancer trial of local treatment⁴⁴ losses to follow up were 1.6% at 5 years. For any patients that are loss to follow-up, sites should be able to obtain primary outcome data from their medical notes.

12.1.2 Non-inferiority Margin

Following consultation with relevant breast cancer consumer groups, lay people, and with breast surgeons and oncologists we have opted for a 2% non-inferiority margin. The consensus across all these groups was that a 2% margin for axillary recurrence is an acceptable trade-off for the increase in arm morbidity associated with axillary treatment.

Axillary recurrence following axillary treatment is estimated to be 2% at 5 years. Therefore, a 2% non-inferiority margin means accepting that 4% of women who have adjuvant therapy alone will have axillary recurrence at 5 years. This is lower than the level recommended by the Association of Breast Surgery, whose guidelines state that axillary recurrence at 5 years should be less than 5%⁴³. This margin appears to be acceptable to women with early breast cancer, as demonstrated in a trial of alternative treatments¹⁷ that used a 2% non-inferiority margin based on axillary recurrence and successfully recruited 4806 participants.

12.2 Statistical Methods

Trial Feasibility Phase

The trial has an internal feasibility phase where an assessment will be made after 12 months recruitment as to whether to continue with the study. This assessment will be made by the independent Trial Steering Committee (TSC), in consultation with the Data Monitoring Committee. The assessment will be against criteria for recruitment, compliance with the intervention, and retention agreed a priori with these committees. The TSC will then recommend to the funder and the sponsor whether the trial should continue.

Recruitment to the trial will continue whilst the feasibility assessment is being conducted.

Stopping guidelines shall be defined in consultation with the DMC.

12.3 Statistical Analysis

Analysis will be at the end of the five years follow up of the last recruited participant. The detailed Statistical Analysis Plan will be developed by the Trial Statisticians in consultation with the Trial Management Group, and agreed with the TSC before database lock and unblinding of the data.

Recruitment at each site will be summarised, along with the main reasons why eligible women were not recruited. For women recruited to the trial, baseline characteristics in the two allocated groups will be described, and compliance with the allocated intervention documented.

The primary analysis will be per protocol for the primary and secondary endpoints. Intention to treat (ITT) analysis will not be performed as the primary analysis because if there are individuals who default from the allocated intervention, the observed difference between the two groups may be reduced. Therefore, an ITT analysis may wrongly declare non-inferiority and is not appropriate for this study.

The primary endpoint is whether axillary recurrence has occurred by 5 years or not. A binary endpoint of this form is judged to be of greatest relevance, in this trial.

The primary analysis will assess whether adjuvant therapy alone is non-inferior to adjuvant therapy plus axillary treatment based on the primary endpoint of axillary recurrence at 5 years. Non-inferiority of adjuvant therapy alone over adjuvant therapy plus axillary treatment will be accepted if the upper boundary of the Miettinen-Nurminen two-sided 95% confidence interval around the estimated difference in axillary recurrence rate lies is no greater than 2%. Secondary intention to treat and CACE (Complier Average Causal Effect) analyses will be performed for the primary and secondary

endpoints. All outcome measures will be compared between the two groups by standard methods, including summary statistics and confidence intervals for measures of effect size. Proportional hazards regression analysis will be performed for time-to-event outcomes such as time to axillary recurrence, axillary recurrence free, disease free and overall survival up to 5 years. Appropriate adjustments will be conducted in case of violations of assumptions of proportionality of hazards and time independence. Other secondary binary outcomes will be compared between the two groups by estimating the differences and ratios of proportions and 95% confidence intervals.

Competing risk methodology will be applied for estimating the disease free survival and the axillary recurrence free survival where death is a competing event.

Subgroup analyses

The following pre-specified subgroup analyses based on criteria at trial entry shall be performed:

1. Number of sentinel node macrometastases: 1, 2
2. Age : <50 years, ≥50 years
3. Breast surgery: mastectomy, breast conserving surgery
4. Estrogen receptor (ER) status: positive,negative
5. Tumour grade: grades 1 or 2, grade 3
6. Sentinel node assessment technique: OSNA, non-OSNA
7. Extranodal invasion: present, absent

We will perform subgroup analyses for the outcomes: axillary recurrence and arm morbidity (measured by LBCQ (2 questions), QuickDash and FACT B+4 arm morbidity subscale). We will describe the outcomes for each treatment by factor subgroup, and present between-group effects and 95% confidence intervals for each factor level. We will include interaction terms in the appropriate regression models, and present estimated interaction effects and 95% confidence intervals for all subgroup analyses conducted. Due to the limited power available to detect all except large interactions, these subgroup analyses will be regarded as exploratory.

Quality of Life

Quality of life outcomes will be analysed according to per protocol analysis. Any missing data and / or imputation will follow standardised rules as defined by the relevant manual for the relevant questionnaires. Any changes in state anxiety over time will be examined using analysis of covariance (ANCOVA) with trait anxiety and baseline state as covariates. Mean change scores for FACT B+4, TOI and subscales will be compared between groups using unpaired t tests at each time point and using ANCOVA over time adjusting for the baseline dependant variable.

12.4 Analysis Population

Participants who receive their allocated treatment will be included in the primary per protocol analysis. Secondary analysis will be intention to treat and CACE including all participants randomised to each treatment group regardless of their received intervention.

12.5 Missing Data

Every effort will be made to minimise the occurrence of missing data. We will examine the plausibility that primary outcome data are missing at random (MAR) and multiple imputation techniques will be used to handle missing values as appropriate. Sensitivity analyses will be considered for testing the robustness of the MAR assumption. A complete case analysis will be performed for secondary outcomes with less than 1% missing data, otherwise techniques similar to primary outcome will be applied.

13 HEALTH ECONOMIC ANALYSIS

The economic evaluation will be carried out from the perspective of the NHS over a three year time horizon. A three year time horizon was adopted for the economic evaluation as it was anticipated, at three years following the initial biopsy, any health-related quality of life (HRQoL) differences between the intervention and standard care participants which might have been present earlier will have disappeared. Costs will be based upon the costs of the randomised interventions received and on the use of subsequent care and services.

For each participant, all clinical events (i) relevant to management and consequences following the initial sentinel node biopsy (adjuvant therapy alone or adjuvant therapy plus axillary treatment) (ii) up to the end of follow-up (iii) which occur in the hospital setting, will be recorded. These include primary treatments, re-admissions for complications or recurrences and associated adverse events. Events deemed unrelated to the consequences of the sentinel node biopsy or axillary treatment will not be included. At 6, 12, 24, 36, 48 and 60 months follow-up, participants will be asked to recall the extent to which they accessed primary care (e.g. GP) or other non-hospital services (e.g. physio visits) for their breast cancer. Information relating to healthcare resource use will be collected via the eCRF. Costs will be estimated using routine sources (e.g. NHS reference costs, PSSRU). On the basis of each participant's record and the costs collected from routine sources, a total hospital cost and a total non-hospital cost will be derived for each participant for the duration of follow-up. These total costs will be combined to estimate the total cost per participant from which the total average cost per randomised group will be calculated. The breast and sentinel node biopsy procedure, systemic treatment, radiotherapy to breast or chest wall will not be included in the determination of costs, because all participants will undergo this.

- 1) **Cost-effectiveness analysis** based on the incremental cost per reduction in axillary recurrence. Mean costs for each randomised arm will be calculated as will mean axillary recurrence. In the cost-effectiveness analysis these will then be presented as point estimates of mean incremental costs and effects (reduced axillary recurrences) and the incremental cost per reduction in axillary recurrence.
- 2) **Cost-utility analysis** based on the incremental cost per QALY gained. QALYs will be based on the area under the curve approach⁴⁵ using responses to the EQ-5D-5L completed at scheduled time points; baseline, 3, 6, 12, 24, and 36 months post randomisation. The total average QALYs accruing in the intervention and standard care groups will be analysed for significant differences, following allowances for mortality (if occurring). Both mean cost and QALYs will be presented for each randomised group and incremental mean costs and QALY calculated along with the incremental cost per QALY gained.

For both the cost-effectiveness and cost-utility analyses the results will be presented as point estimates of mean incremental costs and effects as well as in stochastic analyses plots of cost and

effects and cost-effectiveness acceptability curves. Both deterministic and stochastic sensitivity analyses will be adopted to address any uncertainty in our estimation of costs, effects and cost-effectiveness. Costs and effects will be discounted at UK recommended rates⁴⁶ as they are being estimated over a three year time horizon.

14 REGULATORY AND PARTICIPANT INFORMED CONSENT

14.1 Research Ethics Approval

The Chief Investigator (CI), or their delegate, will obtain approval from the Research Ethics Committee (REC), which will be submitted for Site Specific Assessment (SSA) at each participating NHS Trust. The Principal Investigator (PI) will require a R&D trust approval letter before approaching potential participants for the study.

The CI will provide reports to REC at the intervals stipulated in the REC guidelines.

14.1.1 Protocol Amendments

If it is proposed to make a substantial amendment to the protocol, the CI shall submit a notice of amendment to REC. A substantial amendment is any amendment to the terms of the application for ethical review, or to the protocol or other supporting documentation approved by REC that is likely to affect to a significant degree:

- a) the safety or physical or mental integrity of the trial participants
- b) the scientific value of the trial
- c) the conduct or management of the trial.

The agreement of the sponsor will be sought before submitting the notice of amendment.

A substantial amendment will not be implemented until a favourable ethical opinion has been given by REC, unless the changes to the research are urgent safety measures.

Amendments that are not substantial amendments (“minor amendments”) may be made at any time and do not need to be notified to REC.

14.1.2 Ethical Considerations

The ethical issue is that women randomised to adjuvant therapy alone will not undergo axillary node clearance or axillary radiotherapy and this may be unsettling. Women may be worried that they will potentially have inferior outcomes in terms of cancer recurrence and longevity.

This will be addressed by advising women clearly that we do not know which one is better and both treatments are acceptable. Communication training sessions will be held at participating sites and a participant information DVD will be used to complement the participant information leaflet. The information including randomisation and clinical equipoise will be given in a balanced way. Patient

groups such as Independent Cancer Patients' Voice (ICPV) will explore the interpretation of study information.

14.2 Consent

Potential participants are women whose node is positive at sentinel node biopsy. Based on local hospital practices, there are four pathways for recruitment into the trial. In most hospitals, the results of the biopsy are not available until after the sentinel node biopsy surgery but in a few hospitals (less than 10%), the sentinel node assessment is done whilst the woman remains anaesthetised in the operating room. Women who come back to the clinic to discuss their surgery results will be consented post breast surgery (Pathway 1). At hospitals where sentinel node analysis is performed during the surgery, women will usually be approached and consented for the trial prior to their breast surgery (Pathway 2). However, women at these hospitals may follow Pathway 1. Participants in pathway 2 will be made aware that they will be randomised only if the sentinel node is positive. Women may be consented during or after they have received adjuvant chemotherapy as long as they have not received any axillary treatment (Pathway 3). Women who have had a sentinel node biopsy prior to neoadjuvant therapy can be randomised before, during or after their neoadjuvant therapy as long as they have not received any axillary treatment (Pathway 4).

Consent will be taken by appropriately trained healthcare professionals in the research team.

If the woman does not wish to participate, she will not be required to give a reason. Clinical care will not be influenced by whether or not she agrees to participate. A local site screening log will be maintained recording all potentially eligible women not approached to give consent, women who decline participation, and reasons for non-participation (if available).

The original consent form (CF) will be kept in the Site Study File and a copy will be given to the participant and a copy will be held in the participant's medical notes. With the participant's explicit consent a copy of the signed consent form will be sent to the Nottingham Clinical Trials Unit for central monitoring purposes.

14.3 Confidentiality

Information about participants will be kept confidential and managed according to the requirements of the Data Protection Act, NHS Caldicott Guardian, Research Governance Framework for Health and Social Care, conditions of REC approval and NHS information governance policy.

A unique identification number will be automatically attributed to each participant randomised in the study.

Participants' addresses and telephone numbers will be captured in a separate secure database and will be used by researchers at NCTU and SHORE-C (Sussex University). SHORE-C will be posting questionnaires to the participants' homes and may contact them by telephone in case of queries. Participant contact details will be confidentially destroyed after the last questionnaire has been received and the queries have been resolved.

14.4 Access to Data

Data collection forms will be treated as confidential documents, and held securely. The PI at each site will make a separate confidential record of the participant to permit identification of all participants enrolled in the study. This information will be stored securely and separately to the anonymised study data.

Access to all data at site will be restricted to personnel approved by the local Principal Investigator, and recorded on a delegation log. Access will also be given to the sponsor, REC and NCTU representatives.

15 QUALITY ASSURANCE

This trial will be conducted in accordance with the current approved protocol, ICH GCP, Research Governance Framework for Health and Social Care, principles of the Declaration of Helsinki, all applicable Standard Operating Procedures (SOPs) and all local Trust policies that impact on the conduct of research and any subsequent amendments.

15.1 Site Start-up and Training

Before trial recruitment begins, a meeting with the Principal Investigators and the local research team will be organised to discuss protocol issues, trial interventions and related clinical issues, data collection issues, and trial procedures.

Each hospital will have a trial Site Initiation Visit (SIV), combining training in the trial procedures from NCTU and QoL metrics by SHORE-C, before their first participant is recruited. SIVs will be coordinated with SHORE-C and NCTU attending the meeting.

15.2 Monitoring

Trial monitoring will be in accordance with the monitoring plan and will be a combination of central monitoring and "triggered" site visits when required. Central monitoring will be used to monitor patterns of recruitment at sites, reasons for non-recruitment of potentially eligible participants, characteristics of participants recruited, time of recruitment, etc. It will also be used to assess compliance with the protocol, which may include checking compliance with the trial interventions.

Based on assessment of data processing and central monitoring, the TMG will decide if any further action needs to be taken.

When site visits are performed, a random sample of participants will have their data monitored at source (Source Document Verification). Any major discrepancies or concerns at a site visit will trigger a more extensive audit of trial data at that site.

The investigator at a site will permit direct access to source data and study related documentation for study-related monitoring, audits and inspections by the REC, the sponsor and the Nottingham Clinical Trials Unit (NCTU). This study will be monitored by the NCTU according to a monitoring plan, agreed with the sponsor, and the NCTU procedures for monitoring. In line with the responsibilities set out in the Research Governance Framework, the investigator will ensure that the trial manager, sponsor and/or REC representatives are given access to all trial-related documents, including participant medical notes to enable source data verification and trial related facilities.

15.3 Radiotherapy Quality Assurance

There is worldwide evidence to justify the inclusion of radiotherapy quality assurance (RTQA) programmes as an integral part of clinical trial protocols⁴⁷⁻⁵⁰. They serve to improve protocol compliance and, in a multi-centre setting, minimise variations ensuring clinical trial outcomes reflect differences in randomisation schedules rather than departures from study protocol.

The NCRI Radiotherapy Trials QA (RTTQA) group will be responsible for implementing and coordinating the Radiotherapy Quality Assurance programme. The QA programme is outlined below:

Pre-trial QA - Prior to site activation, the following must be completed:

- Facility questionnaire – Details of treatment technique, immobilisation, verification and dosimetry will be recorded.
- Control case - All centres must submit a test participant case to the RTTQA group, for review, demonstrating planned radiotherapy treatment to the breast/chest wall, axilla and supraclavicular fossa. The outlining, planning and treatment of this participant case should reflect how the centre intends to plan and treat all participants recruited to POSNOC.

On-trial QA – Ongoing QA requirements

- Retrospective individual case reviews –All centres must export data of their first ten radiotherapy plans to the RTTQA group. Data will include participant history, CT data, structure set, plan and dose files. The retrospective review process should include at least three plans with radiotherapy to the nodal region.
- Ongoing data collection - All radiotherapy plans will be exported to the RTTQA group, regardless of which study allocation. A random sample of plans will be reviewed from each site during trial recruitment.

The sites will be required to adhere to the specific radiotherapy planning and delivery guidelines when treating participants in the study. For data anonymisation and export please see section 'Data Collection'.

The NCRI Radiotherapy Trials QA (RTTQA) group will promote POSNOC radiotherapy planning and delivery guidelines compliance for the duration of trial recruitment by systematically reviewing a sample of radiotherapy plans submitted by centres.

RTTQA will monitor compliance to the RT planning and delivering guidelines. Any RT guidelines deviations, non-compliance or areas of concern will be recorded and if required, discussed with the TMG and individual centres. The RTTQA has the responsibility for collection and storage of

radiotherapy data related to the RT QA programme. There will be a named RT QA contact for the POSNOC trial.

16 TRIAL MANAGEMENT

16.1 Trial Management Group

Day-to-day management of the study will be the responsibility of the Trial Management Group (TMG), which will meet every one to two months during the recruitment phase and every six months thereafter.

The TMG will review recruitment, retention, compliance and data quality to ensure efficient study conduct according to the research timeline. They will report to the independent Trial Steering Committee (TSC).

16.2 Trial Steering Committee

The Trial Steering Committee (TSC) will provide independent oversight of the study. They will meet (in person or by telephone conference) prior to commencement of the study, and then at regular intervals until completion (at least annually). Specific tasks of the TSC are:

- to approve the trial protocol
- to approve necessary changes to the protocol based on considerations of feasibility and practicability
- agree trial stopping rules for the feasibility and the main trial
- to receive reports from the Data Monitoring Committee
- to resolve problems brought to it by the co-ordinating centre and TMG
- to ensure publication of the trial results

16.3 Data Monitoring Committee

A separate and independent Data Monitoring Committee (DMC) will be convened. It is anticipated that the members will meet once to agree terms of reference and on at least three further occasions to monitor accumulating data and oversee safety issues. This Committee will be independent of the study organisers and the TSC. During the period of recruitment to the study, data reports will be supplied, in strict confidence to the DMC, together with any other reports that the committee may request. This may include reports of data from other comparable trials. In the light of these data reports, the DMC will advise the Steering Committee if, in its view, there are any ethical or safety issues that may necessitate modification to the protocol or closure of the trial.

The TSC, TMG, clinical collaborators and Coordinating Centre staff (except those who supply the confidential analyses) will not receive copies of these data reports.

The frequency of interim reports will depend on the judgement of the Chairman and other independent DMC members. We anticipate that there might be two interim reports and one final report.

16.4 Indemnity

NHS indemnity scheme will apply in the event of a claim by, or on behalf of, participants for negligent harm. There will be no special arrangements for non-negligent harm but the normal NHS complaints mechanism will be available to all participants.

16.5 Public and Patient Involvement

As part of the development of the project and protocol, women from the Derby Breast Cancer Support Group, Bosom Buddies (Guildford Breast Cancer Support Group) and Independent Cancer Patients' Voice (ICPV) have commented at all stages on the design and acceptability of the study. The Breast Cancer Care and Lymphoedema Support Network have had the opportunity to contribute to this protocol.

Independent Cancer Patients' Voice is an independent patient advocate group (<http://independentcancerpatientsvoice.org.uk/>). Two members of ICPV will provide a patient perspective throughout the duration of the study.

16.6 Security and Back-up of Data

Computer held data including the study database will be held securely and password protected. All data will be stored on a secure dedicated server. Access will be restricted by user identifiers and passwords (encrypted using a one way encryption method). Electronic data will be backed up every 24 hours.

16.7 Archiving

The CI is the "Custodian" of the data. Data and all appropriate study documentation will be archived for 5 years after the end of the study, including the final report and publication in peer reviewed journal. The trial master file and study documents held by the CI and NCTU on behalf of the sponsor will be archived in secure archive facilities by Derby Teaching Hospitals NHS Foundation Trust. This archive will include all study databases and associated meta-data encryption codes.

Quality of life questionnaires and the accompanying database will be securely archived by SHORE-C.

Exported radiotherapy data will be securely archived by the RTTQA group.

16.8 Dissemination policy

16.8.1 Dissemination plan

The dissemination of the study results will be via a study report and research papers for publication in peer reviewed journals, and presentation at relevant conferences. Reporting will be in compliance with CONSORT recommendations. Publication of the results will be based on outcomes at 5 years following the last recruited patient. No interim publication of results is planned, any unscheduled interim publication would require approval of the TSC.

A summary of the results will be made available to participants through a newsletter (unless they state they do not wish to receive this), and will also be publicised through Independent Cancer Patients' Voice, Cancer Research UK, Breast Cancer Care, Breakthrough Breast Cancer and Lymphoedema Support Network.

16.8.2 Policy for Publication and Authorship

The publication and authorship policy shall be agreed with the collaborators. The first author will be the CI of the study. Authorship will be named authors on behalf of a collaborative group, the named authorship is for those who have made a significant contribution. Additional authors will be those who have contributed to the overall success of the study.

Outcomes by treatment group will not be available for publication before the first results paper has been published.

Reference List

- (1) Cancer Research UK. <http://www.cancerresearchuk.org/cancer-info/cancerstats/types/breast/> 2013.
- (2) Veronesi U, Paganelli G, Viale G, Luini A, Zurrada S, Galimberti V et al. A randomized comparison of sentinel-node biopsy with routine axillary dissection in breast cancer. *N Engl J Med* 2003; **349**(6):546-553.
- (3) Mansel RE, Fallowfield L, Kissin M, Goyal A, Newcombe RG, Dixon JM et al. Randomized multicenter trial of sentinel node biopsy versus standard axillary treatment in operable breast cancer: the ALMANAC Trial. *J Natl Cancer Inst* 2006; **98**(9):599-609.
- (4) Krag DN, Anderson SJ, Julian TB, Brown AM, Harlow SP, Costantino JP et al. Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer: overall survival findings from the NSABP B-32 randomised phase 3 trial. *Lancet Oncol* 2010; **11**(10):927-933.
- (5) Goyal A, Newcombe RG, Chhabra A, Mansel RE. Factors affecting failed localisation and false-negative rates of sentinel node biopsy in breast cancer--results of the ALMANAC validation phase. *Breast Cancer Res Treat* 2006; **99**(2):203-208.
- (6) Gill G. Sentinel-lymph-node-based management or routine axillary clearance? One-year outcomes of sentinel node biopsy versus axillary clearance (SNAC): a randomized controlled surgical trial. *Ann Surg Oncol* 2009; **16**(2):266-275.
- (7) Mansel RE, MacNeill F, Horgan K, Goyal A, Britten A, Townson J et al. Results of a national training programme in sentinel lymph node biopsy for breast cancer. *Br J Surg* 2013.
- (8) Breast cancer (early & locally advanced): diagnosis and treatment. <http://guidance.nice.org/uk/CG80> 2009.
- (9) Aitken RJ, Gaze MN, Rodger A, Chetty U, Forrest AP. Arm morbidity within a trial of mastectomy and either nodal sample with selective radiotherapy or axillary clearance. *Br J Surg* 1989; **76**(6):568-571.
- (10) Arriagada R, Mouriessse H, Rezvani A, Sarrazin D, Clark RM, DeBoer G et al. Radiotherapy alone in breast cancer. Analysis of tumor and lymph node radiation doses and treatment-related complications. The experience of the Gustave-Roussy Institute and the Princess Margaret Hospital. *Radiother Oncol* 1993; **27**(1):1-6.

-
- (11) Chetty U, Jack W, Prescott RJ, Tyler C, Rodger A. Management of the axilla in operable breast cancer treated by breast conservation: a randomized clinical trial. Edinburgh Breast Unit. *Br J Surg* 2000; **87**(2):163-169.
 - (12) Hack TF, Cohen L, Katz J, Robson LS, Goss P. Physical and psychological morbidity after axillary lymph node dissection for breast cancer. *J Clin Oncol* 1999; **17**(1):143-149.
 - (13) Hoebbers FJ, Borger JH, Hart AA, Peterse JL, Th EJ, Lebesque JV. Primary axillary radiotherapy as axillary treatment in breast-conserving therapy for patients with breast carcinoma and clinically negative axillary lymph nodes. *Cancer* 2000; **88**(7):1633-1642.
 - (14) Roses DF, Brooks AD, Harris MN, Shapiro RL, Mitnick J. Complications of level I and II axillary dissection in the treatment of carcinoma of the breast. *Ann Surg* 1999; **230**(2):194-201.
 - (15) Glechner A, Wockel A, Gartlehner G, Thaler K, Strobelberger M, Griebler U et al. Sentinel lymph node dissection only versus complete axillary lymph node dissection in early invasive breast cancer: A systematic review and meta-analysis. *Eur J Cancer* 2013; **49**(4):812-825.
 - (16) Straver ME, Meijnen P, van Tienhoven G, van de Velde CJ, Mansel RE, Bogaerts J et al. Role of axillary clearance after a tumor-positive sentinel node in the administration of adjuvant therapy in early breast cancer. *J Clin Oncol* 2010; **28**(5):731-737.
 - (17) Rutgers EJ, Donker M, Straver ME, Meijnen P, van de Velde CJ, Mansel R et al. Radiotherapy or surgery of the axilla after a positive sentinel node in breast cancer patients: Final analysis of the EORTC AMAROS trial (10981/22023). *J Clin Oncol* 2013; **31**:suppl; abstr LBA1001.
 - (18) Damera A, Evans AJ, Cornford EJ, Wilson AR, Burrell HC, James JJ et al. Diagnosis of axillary nodal metastases by ultrasound-guided core biopsy in primary operable breast cancer. *Br J Cancer* 2003; **89**(7):1310-1313.
 - (19) Fisher B, Jeong JH, Anderson S, Bryant J, Fisher ER, Wolmark N. Twenty-five-year follow-up of a randomized trial comparing radical mastectomy, total mastectomy, and total mastectomy followed by irradiation. *N Engl J Med* 2002; **347**(8):567-575.
 - (20) Louis-Sylvestre C, Clough K, Asselain B, Vilcoq JR, Salmon RJ, Campana F et al. Axillary treatment in conservative management of operable breast cancer: dissection or radiotherapy? Results of a randomized study with 15 years of follow-up. *J Clin Oncol* 2004; **22**(1):97-101.

-
- (21) Haffty BG, Ward B, Pathare P, Salem R, McKhann C, Beinfield M et al. Reappraisal of the role of axillary lymph node dissection in the conservative treatment of breast cancer. *J Clin Oncol* 1997; **15**(2):691-700.
- (22) Renolleau C, Merviel P, Clough KB, Asselain B, Campana F, Durand JC. Isolated axillary recurrences after conservative treatment of breast cancer. *Eur J Cancer* 1996; **32A**(4):617-621.
- (23) Recht A, Pierce SM, Abner A, Vicini F, Osteen RT, Love SM et al. Regional nodal failure after conservative surgery and radiotherapy for early-stage breast carcinoma. *J Clin Oncol* 1991; **9**(6):988-996.
- (24) Buchholz TA, Tucker SL, Erwin J, Mathur D, Strom EA, McNeese MD et al. Impact of systemic treatment on local control for patients with lymph node-negative breast cancer treated with breast-conservation therapy. *J Clin Oncol* 2001; **19**(8):2240-2246.
- (25) Khonji NI, Edwards R, Sweetland HM, Monypenny IJ, Mansel RE, Webster DJ. Incidence of nodes in completion mastectomy specimens following breast conservation and axillary clearance. *Br J Surg* 2002; **89**(10):1294-1295.
- (26) Veronesi U, Orecchia R, Zurrada S, Galimberti V, Luini A, Veronesi P et al. Avoiding axillary dissection in breast cancer surgery: a randomized trial to assess the role of axillary radiotherapy. *Ann Oncol* 2005; **16**(3):383-388.
- (27) Giuliano AE, McCall L, Beitsch P, Whitworth PW, Blumencranz P, Leitch AM et al. Locoregional recurrence after sentinel lymph node dissection with or without axillary dissection in patients with sentinel lymph node metastases: the American College of Surgeons Oncology Group Z0011 randomized trial. *Ann Surg* 2010; **252**(3):426-432.
- (28) Jagsi R, Chadha M, Moni J, Ballman K, Laurie F, Buchholz TA et al. Radiation Field Design in the ACOSOG Z0011 (Alliance) Trial. *J Clin Oncol* 2014; **32**(32):3600-3606.
- (29) Azghadi S, Daly M, Mayadev J. Practice Patterns of Radiation Field Design for Sentinel Lymph Node-Positive Early-Stage Breast Cancer. *Clin Breast Cancer* 2016; **16**(5):410-417.
- (30) Donker M, van TG, Straver ME, Meijnen P, van de Velde CJ, Mansel RE et al. Radiotherapy or surgery of the axilla after a positive sentinel node in breast cancer (EORTC 10981-22023 AMAROS): a randomised, multicentre, open-label, phase 3 non-inferiority trial. *Lancet Oncol* 2014; **15**(12):1303-1310.
- (31) Hieken TJ, Trull BC, Boughey JC, Jones KN, Reynolds CA, Shah SS et al. Preoperative axillary imaging with percutaneous lymph node biopsy is valuable

in the contemporary management of patients with breast cancer. *Surgery* 2013; **154**(4):831-838.

- (32) Farrell TP, Adams NC, Stenson M, Carroll PA, Griffin M, Connolly EM et al. The Z0011 Trial: Is this the end of axillary ultrasound in the pre-operative assessment of breast cancer patients? *Eur Radiol* 2015; **25**(9):2682-2687.
- (33) Lloyd P, Theophilidou E, Newcombe RG, Pugh L, Goyal A. Axillary tumour burden in women with a fine-needle aspiration/core biopsy-proven positive node on ultrasonography compared with women with a positive sentinel node. *Br J Surg* 2017; **104**(13):1811-1815.
- (34) van Wely BJ, de Wilt JH, Francissen C, Teerenstra S, Strobbe LJ. Meta-analysis of ultrasound-guided biopsy of suspicious axillary lymph nodes in the selection of patients with extensive axillary tumour burden in breast cancer. *Br J Surg* 2015; **102**(3):159-168.
- (35) Verheuel NC, van dH, I, Ooms HW, Voogd AC, Roumen RM. The role of ultrasound-guided lymph node biopsy in axillary staging of invasive breast cancer in the post-ACOSOG Z0011 trial era. *Ann Surg Oncol* 2015; **22**(2):409-415.
- (36) Puri S, Sharma N, Newcombe RG, Kaushik M, Al-Attar M, Pascaline S et al. Axillary tumour burden in women with one abnormal node on ultrasound scan compared to women with multiple abnormal nodes. *Clin Radiol* 2017;in press.
- (37) Armer JM, Radina ME, Porock D, Culbertson SD. Predicting breast cancer-related lymphedema using self-reported symptoms. *Nurs Res* 2003; **52**(6):370-379.
- (38) Kennedy CA, Beaton DE, Smith P, Van ED, Tang K, Inrig T et al. Measurement properties of the QuickDASH (Disabilities of the Arm, Shoulder and Hand) outcome measure and cross-cultural adaptations of the QuickDASH: a systematic review. *Qual Life Res* 2013.
- (39) Coster S, Poole K, Fallowfield LJ. The validation of a quality of life scale to assess the impact of arm morbidity in breast cancer patients post-operatively. *Breast Cancer Res Treat* 2001; **68**(3):273-282.
- (40) Eton DT, Cella D, Yost KJ, Yount SE, Peterman AH, Neuberg DS et al. A combination of distribution- and anchor-based approaches determined minimally important differences (MIDs) for four endpoints in a breast cancer scale. *J Clin Epidemiol* 2004; **57**(9):898-910.
- (41) Spielberger C. The Manual for the State-Trait Inventory. *Palo Alto, CA: Consulting Psychologists Press Inc* 1983.

-
- (42) Sanghani M, Balk EM, Cady B. Impact of axillary lymph node dissection on breast cancer outcome in clinically node negative patients: a systematic review and meta-analysis. *Cancer* 2009; **115**(8):1613-1620.
- (43) Association of Breast Surgery at BASO. Surgical guidelines for the management of breast cancer. *European Journal of Surgical Oncology (EJSO)* 2009; **35**, **Supplement 1**(0):S1-S22.
- (44) Bentzen SM, Agrawal RK, Aird EG, Barrett JM, Barrett-Lee PJ, Bentzen SM et al. The UK Standardisation of Breast Radiotherapy (START) Trial B of radiotherapy hypofractionation for treatment of early breast cancer: a randomised trial. *Lancet* 2008; **371**(9618):1098-1107.
- (45) Matthews JN, Altman DG, Campbell MJ, Royston P. Analysis of serial measurements in medical research. *BMJ* 1990; **300**(6719):230-235.
- (46) National Institute for Health and Care Excellence (NICE): **Guide to the methods of technology appraisal 2013**. <https://www.nice.org.uk/article/pmg9> 2013.
- (47) Abrams RA, Winter KA, Regine WF, Safran H, Hoffman JP, Lustig R et al. Failure to adhere to protocol specified radiation therapy guidelines was associated with decreased survival in RTOG 9704--a phase III trial of adjuvant chemotherapy and chemoradiotherapy for patients with resected adenocarcinoma of the pancreas. *Int J Radiat Oncol Biol Phys* 2012; **82**(2):809-816.
- (48) Weber DC, Poortmans PM, Hurkmans CW, Aird E, Gulyban A, Fairchild A. Quality assurance for prospective EORTC radiation oncology trials: the challenges of advanced technology in a multicenter international setting. *Radiother Oncol* 2011; **100**(1):150-156.
- (49) Peters LJ, O'Sullivan B, Giralt J, Fitzgerald TJ, Trotti A, Bernier J et al. Critical impact of radiotherapy protocol compliance and quality in the treatment of advanced head and neck cancer: results from TROG 02.02. *J Clin Oncol* 2010; **28**(18):2996-3001.
- (50) Purdy JA. Quality assurance issues in conducting multi-institutional advanced technology clinical trials. *Int J Radiat Oncol Biol Phys* 2008; **71**(1 Suppl):S66-S70.

Appendices

1. Glossary of abbreviations / acronyms

AE	Adverse Event
ANC	Axillary Node Clearance
ANZ	Australia and New Zealand
ART	Axillary Radiotherapy
BCS	Breast-conserving surgery
CACE	Complier Average Causal Effect
CF	Consent Form
CI	Chief Investigator
CONSORT	Consolidated Standards of Reporting Trials
CRF	Case Record Form
DMC	Data Monitoring Committee
EORTC	European Organisation for Research and Treatment of Cancer
ER	Estrogen receptor
FACT B+4	Functional Assessment of Cancer Therapy-Breast+4
GCP	Good Clinical Practice
HER2	Human epidermal growth factor receptor 2
HRQL	Health-Related Quality of Life
HSCIC	Health and Social Care Information Centre
HTA	Health Technology Assessment
ICER	Incremental Cost Effectiveness Ratio
ITT	Intention To Treat
ICPV	Independent Cancer Patients' Voice
LBCQ	Lymphoedema and Breast Cancer Questionnaire
MAR	Missing At Random
NCTU	Nottingham Clinical Trials Unit
NHS	National Health Service
NICE	National Institute of Clinical Excellence
NIHR	National Institute of Health Research
PI	Principal Investigator at a local centre
PIS	Participant Information Sheet
QoL	Quality of Life
QALYs	Quality Adjusted Life-Years
RCT	Randomised Control Trial
REC	Research Ethics Committee
RTTQA	NCRI Radiotherapy Trials QA
R&D	Research and Development department
SAE	Serious Adverse Event
SAP	Statistical Analysis Plan
SCF	supra clavicular fossa
SHORE-C	Sussex Health Outcomes Research & Education in Cancer
SIV	Site Initiation Visit
SN	Sentinel Node
SNB	Sentinel Node Biopsy
SOP	Standard Operating Procedure
SSA	Site Specific Assessment
TMG	Trial Management Group
TOI	Trial Outcome Index
TSC	Trial Steering Committee